

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A23L 1/105, C12P 19/14, 19/22</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/07628 (43) International Publication Date: 23 March 1995 (23.03.95)</p>
<p>(21) International Application Number: PCT/SE94/00857 (22) International Filing Date: 14 September 1994 (14.09.94) (30) Priority Data: 9302996-5 15 September 1993 (15.09.93) SE (71)(72) Applicants and Inventors: LINDAHL, Lennart [SE/SE]; Uardavägen 16 D, S-224 71 Lund (SE). AHLDEN, Inger [SE/SE]; Tullgatan 8 A, S-223 54 Lund (SE). ÖSTE, Rickard [SE/SE]; Kollegievägen 91, S-224 73 Lund (SE). SJÖHOLM, Ingegerd [SE/SE]; Nöbbelövs Kyrkoväg 55, S-226 53 Lund (SE). (74) Agent: AWAPATENT AB; P.O. Box 5117, S-200 71 Malmö (SE).</p>		<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD). Published With international search report. In English translation (filed in Swedish).</p>

(54) Title: HOMOGENEOUS AND STABLE CEREAL SUSPENSION

(57) Abstract

A homogeneous and stable cereal suspension having the taste and aroma of natural oats, as well as a method for the preparation thereof, is described. The cereal suspension contains intact β -glucans from the starting material and is prepared by A) dry- or wet-grinding rolled oats or otherwise heat- and water-treated oats to meal, B) suspending the oatmeal in water, if the meal has been produced by dry grinding, C) optionally centrifuging or decanting the suspension in order to remove coarse fibre particles, D) treating the suspension with β -amylase, which specifically generates maltose units and has no glucanase and proteinase effect, to a viscosity of 3-0.1 Pas in the shear rate range of 10-100 s⁻¹, E) treating the suspension with α -amylase, which specifically generates maltose units and has no glucanase and proteinase effect, to a viscosity of < 0.5 Pas in the shear rate range of 10-100 s⁻¹, F) preferably homogenising the enzyme-treated suspension, and G) subjecting the suspension to UHT treatment (UHT = Ultra High Temperature) in order to obtain a sterile product while inactivating the enzymes added.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HOMOGENEOUS AND STABLE CEREAL SUSPENSION

This invention relates to a homogeneous and stable cereal suspension tasting and smelling like natural oats, as well as a method for the preparation thereof.

In many respects, oats is different from other cereals. Thus, it
5 has higher protein and fat contents as well as a lower carbohydrate content than comparable cereals.

In recent years, there has been a growing interest for food made from oats. The main reason for this is that oat fibres have been found to have a wholesome effect by lowering the serum cholesterol level.
10 Another reason is that oats contains protein of high food value as well as a considerable proportion of polyunsaturated fats. In addition, oats contains a lot of essential amino acids and minerals.

A great advantage of oats is that the whole grain can be used for making various products, once the hull has been removed. In oats,
15 the most nutritious substances are distributed fairly evenly in the whole grain. In other cereals, the nutritious substances are frequently concentrated in some parts of the grain.

Oats is mostly used for making porridge oats and oatmeal to be added to gruel products and baby food. Oats in various forms is also
20 added to bread, cakes, sweets, bread or cake mixes, health foods, and so forth. The functional properties of oats come in handy in such products as sauces, soups and minced meat.

US Patent Specification 4,996,063 (G.F. Inglett) discloses the preparation of water-soluble dietary-fibre compositions by treating
25 ground oat products with α -amylases. The α -amylase then serves to thin the oat starch, and any α -amylases may thus be used. The thus-produced pulverulent dietary-fibre compositions, which are colourless and lack natural aromatics, are used as additives in food. These prior-art products not only lack undesirable aromatics but are
30 also deprived of the agreeable flavourings and aromatics found in natural oats. These products are used as fat substitutes.

The object of the present invention is to provide a milky product based on oats.

Since it should be possible to drink the milky product as it is, the suspension has to remain stable and homogeneous for a long time and have an appealing taste and aroma. Furthermore, the product should not contain any foreign additives, such as stabilisers and artificial flavourings and aromatics.

According to the invention, this object is achieved by a homogeneous and stable cereal suspension which has the taste and aroma of natural oats and contains intact β -glucans from the starting material and which has been prepared by

- 10 A) dry- or wet-grinding rolled oats or otherwise heat- and water-treated oats to meal,
- B) suspending the oatmeal in water, if the meal has been produced by dry grinding,
- 15 C) optionally centrifuging or decanting the suspension in order to remove coarse fibre particles,
- D) treating the suspension with β -amylase, which specifically generates maltose units and has no glucanase and proteinase effect, to a viscosity of 3-0.1 Pas in the shear rate range of 10-100 s⁻¹,
- 20 E) treating the suspension with α -amylase, which specifically generates maltose units and has no glucanase and proteinase effect, to a viscosity of < 0.5 Pas in the shear rate range of 10-100 s⁻¹,
- F) preferably homogenising the enzyme-treated suspension, and
- 25 G) subjecting the suspension to UHT treatment (UHT = Ultra High Temperature) in order to obtain a sterile product while inactivating the enzymes added.

The invention further comprises a method for preparing a homogeneous and stable cereal suspension that contains intact β -glucans from the starting material and has the taste and aroma of natural oats. The inventive method comprises the steps A)-G) indicated above.

The cereal suspension according to the invention can be used as an alternative to milk. Apart from the above-mentioned nutritious qualities of oats, the product should possess such properties as to be suitable for lactose-intolerant people.

35 The inventive product may be used as basis of or additive to ice-cream, gruel, yoghurt and milkshake, or as health drink or snack between meals.

The product is homogeneous and stable without any additives in the form of foreign stabilisers, since it contains intact β -glucans from the starting material employed.

Conveniently, the cereal suspension according to the invention
5 is prepared on the basis of commercially-produced, pregelatinised rolled oats retaining the original taste and aroma of the oats. The rolled oats is ground to oatmeal by total, dry or wet grinding. In dry grinding, the oatmeal is suspended in water, preferably at a temperature of 50-53°C. Also in wet grinding, use is preferably made of water
10 having a temperature of 50-53°C. Especially good results are obtained if the water has been deionised.

Suitably, the slurry or suspension has a weight ratio of meal to water in the range of 1:5-1:8, which corresponds to a dry solids content of 10-15%. The suspension is agitated until the meal has been
15 dissolved and the desired extraction has been achieved. The slurry should have a pH of at least 5.

In order to remove coarse fibre particles, the suspension can then be centrifuged or decanted at 350-450 G and for about 10-15 min.

20 Then, β -amylase and α -amylase are added. The β -amylase should be a β -amylase that specifically generates maltose units and has no glucanase and proteinase effect, preferably 1,4- α -D-glucan maltohydrolase. The α -amylase should be an α -amylase which specifically generates maltose units and has no glucanase and proteinase
25 effect.

The amount of enzyme, the temperature of the slurry, the agitation time and the pH value are optimised after the gelatinisation of starch in order to obtain a product of suitable viscosity after heating. The choice of suitable parameters is influenced by several factors,
30 such as the starting material employed, the enzymes added, and the desired viscosity of the end product.

The treatment with β -amylase is carried out such that the suspension obtains a viscosity of 3-0.1 Pas in the shear rate range of 10-100 s⁻¹. The treatment with α -amylase is carried out such that
35 the suspension obtains a viscosity of < 0.5 Pas in the same shear rate range.

The product is then homogenised, suitably at a temperature of 72-75°C and a pressure of 200-250 bar.

Finally, the product is subjected to UHT treatment (Food Engineering and Dairy Technology, H.G. Kessler, Verlag A. Kessler, 1981, Chapter 6, pp 139-207) in view of sterilising it while inactivating the enzymes added. Conveniently, the end product is packed aseptically.

The maximum dry solids content of the product is in the range of 10-15%. The viscosity at room temperature is below 0.5 Pas. The β -glucan values (based on the dry solids content) were determined with the aid of a so-called Biocon kit from Biocon Pty. Ltd., Australia, as follows:

Cereal product not subjected to enzyme treatment:	6.5% on DS
Cereal product + β -amylase:	5.8% on DS

The invention will now be described in more detail with the aid of two Examples.

Example 1 (no centrifugation)

Immediately before the other steps of the inventive method, steam-processed rolled oats is ground to meal (100% grinding). It is an advantage if the meal is finely ground such that it can pass through a bolter with a mesh of 0.8-1 mm. 1 kg of meal is mixed with 6 l of water while agitated. The water should have a temperature of 50-53°C.

In a first enzyme-treatment step, β -amylase is added to the formed slurry under continuous agitation. In enzyme step 1, the incubation temperature is 53-55°C. Enzyme step 1 proceeds until the viscosity of the slurry has fallen to a value of 3-0.1 Pas in the shear rate range of 10-100 s⁻¹.

Then, α -amylase which specifically generates maltose units is added to the slurry in a second enzyme-treatment step. In enzyme step 2, the incubation temperature is 55-57°C. The incubation with α -amylase proceeds until the slurry has obtained a viscosity of < 0.5 Pas in the shear rate range indicated above.

Then, the slurry is homogenised at a pressure of 200 bar (homogenisation should be carried out in the range of 160-250 bar) and at a temperature of 72-75°C. The slurry is then immediately treated by indirect vapour at a temperature of 137-138°C and for 3-4 s in order to sterilise the product (killing bacteria and spore-

5

forming agents) before it is packed aseptically. At the same time, the enzymes added are totally inactivated.

The resulting oat base has the following composition

	% DS	13.4
5	% protein on DS	2.2
	% fat on DS	0.8
	% fibres, total on DS	0.8
	of which % soluble fibres	0.34
	% insoluble fibres	0.46
10	% starch on DS (incl. * sugars)	7.9
	* sugars, mg/g DS:	-
	Fructose	2.2
	Glucose	11.8
15	Saccharose	316.0
	Maltose	3.0
	Raffinose	7.8
	Maltotriose	

Example 2 (with centrifugation)

- 20 Example 1 was repeated, but with a separation step for removing coarse fibre particles. The separation was produced by centrifugation at 400 G before the first enzyme incubation.

CLAIMS

1. A homogeneous and stable cereal suspension having the
5 taste and aroma of natural oats, c h a r a c t e r i s e d in that it
contains intact β -glucans from the starting material, and that it has
been prepared by
- A) dry- or wet-grinding rolled oats or otherwise heat- and water-treat-
ed oats to meal,
 - 10 B) suspending the oatmeal in water, if the meal has been produced
by dry grinding,
 - C) optionally centrifuging or decanting the suspension in order to
remove coarse fibre particles,
 - D) treating the suspension with β -amylase, which specifically gene-
15 rates maltose units and has no glucanase and proteinase effect,
to a viscosity of 3-0.1 Pas in the shear rate range of 10-100 s⁻¹,
 - E) treating the suspension with α -amylase, which specifically gene-
rates maltose units and has no glucanase and proteinase effect,
to a viscosity of < 0.5 Pas in the shear rate range of 10-100 s⁻¹,
 - 20 F) preferably homogenising the enzyme-treated suspension, and
G) subjecting the suspension to UHT treatment (UHT = Ultra High
Temperature) in order to obtain a sterile product while inactivating
the enzymes added.

2. A method for preparing a homogeneous and stable cereal
25 suspension containing intact β -glucans from the starting material and
having the taste and aroma of natural oats, c h a r a c t e r i s e d
by the steps of
- A) dry- or wet-grinding rolled oats or otherwise heat- and water-treat-
ed oats to meal,
 - 30 B) suspending the oatmeal in water, if the meal has been produced
by dry grinding,
 - C) optionally centrifuging or decanting the suspension in order to
remove coarse fibre particles,
 - D) treating the suspension with β -amylase, which specifically gene-
35 rates maltose units and has no glucanase and proteinase effect,
to a viscosity of 3-0.1 Pas in the shear rate range of 10-100 s⁻¹,

- E) treating the suspension with α -amylase, which specifically generates maltose units and has no glucanase and proteinase effect, to a viscosity of < 0.5 Pas in the shear rate range of $10-100 \text{ s}^{-1}$,
F) preferably homogenising the enzyme-treated suspension, and
5 G) subjecting the suspension to UHT treatment (UHT = Ultra High Temperature) in order to obtain a sterile product while inactivating the enzymes added.

3. A method as set forth in claim 2, characterised by using deionised water in the wet grinding of step A) and the suspension of step B).

10 4. A method as set forth in claim 2 or 3, characterised by suspending the oatmeal to a dry solids content of 10-15% in step B).

5. A method as set forth in one or more of claims 2-4, characterised by using water of a temperature of $50-53^\circ\text{C}$ in the wet grinding of step A) and the suspension of step B).

15 6. A method as set forth in one or more of claims 2-5, characterised by using 1,4- α -D-glucan maltohydrolase as β -amylase in step D).

20 7. A method as set forth in one or more of claims 2-6, characterised by packing the homogeneous and stable cereal product aseptically.

INTERNATIONAL SEARCH REPORT

1

International application No.

PCT/SE 94/00857

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A23L 1/105, C12P 19/14, C12P 19/22
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A23L, A23J, C12N, C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, WPIL, CLAIMS, CHEMICAL ABSTRACT, BIOSIS, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 4996063 (G.F.INGLETT), 26 February 1991 (26.02.91), column 2, line 59 - column 4, line 11 --	1-7
A	US, A, 4377602 (E. CONRAD), 22 March 1983 (22.03.83), column 2, line 21 - line 23; page 35, column 3, line 49 - column 4, line 10 --	1-7
A	Dialog Information Services, file 51, FSTA, Dialog accession no. 00415982, FSTA accession no. 91-09-n0001, Inglett G.E. et al: "Maltodex- trin fat substitute lowers cholesterol", & Food Technology 1991, 45 (6) p104 -- -----	1-7

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

16 December 1994

22-12-1994

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Anneli Jönsson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

26/11/94

International application No.

PCT/SE 94/00857

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4996063	26/02/91	AU-B- 618491	19/12/91
		AU-A- 5933290	17/01/91
		CA-A- 2035472	31/12/90
		DE-T- 431139	24/02/94
		EP-A- 0431139	12/06/91
		ES-B- 2032245	16/12/93
		ES-T- 2051673	01/07/94
		JP-T- 3503367	01/08/91
		US-A- 5082673	21/01/92
		WO-A- 9100027	10/01/91
US-A- 4377602	22/03/83	AT-B- 365420	11/01/82
		AU-B- 522176	20/05/82
		AU-A- 4063078	01/05/80
		BE-A- 871364	15/02/79
		CA-A- 1120775	30/03/82
		CH-A- 644736	31/08/84
		DE-A,C- 2844896	19/04/79
		FR-A,B- 2406665	18/05/79
		GB-A,B- 2007960	31/05/79
		JP-C- 1510048	26/07/89
		JP-A- 54095760	28/07/79
		JP-B- 63063179	06/12/88
		LU-A- 80380	19/03/79
		NL-A- 7810395	20/04/79
		SE-B,C- 429492	12/09/83
		SE-A- 7808870	19/04/79
		US-A- 4282319	04/08/81